

Nonlinear α -Effect in Dynamo Theory

by

George B. Field, Eric G. Blackman, and Hongsong Chou

Received _____; accepted _____

ABSTRACT

The standard two-scale theory of the dynamo coefficient α in incompressible isotropic helical MHD turbulence is extended to include nonlinear effects of $\overline{\mathbf{B}}$, the large-scale magnetic field. We express α in terms of statistical quantities that can be calculated from numerical simulations of the case $\overline{\mathbf{B}} = 0$. For large magnetic Reynolds numbers our formula agrees approximately with that of Kraichnan (1979), but disagrees with that of Cattaneo and Hughes (1996).

1. Introduction

Magnetic fields of galaxies are important in astrophysics and cosmology. In astrophysics, because they enable fast particles to be accelerated and trapped, and affect the dynamics of star formation. In cosmology, because if galactic magnetic fields do not originate in the modern era, they could be relics from the early universe, carrying information about that period.

Since Parker’s (1955) paper on the α - Ω dynamo, and his application of dynamo theory to the Galaxy (1971), most workers have attributed the origin of the magnetic fields of disk galaxies to the operation of an α - Ω turbulent dynamo (Ruzmaikin, Shukurov & Sokoloff 1988). However, the standard theory of the dynamo has always been open to criticism. For example, Piddington (1970, 1972abc, 1975ab) argued that the small-scale magnetic field produced by the small-scale turbulence required by the theory would rapidly grow to equipartition, quenching dynamo action, and this point has been demonstrated numerically by Kulsrud and Anderson (1992).

Recently a series of papers (Vainshtein & Rosner 1991; Cattaneo & Vainshtein 1991; Vainshtein and Cattaneo 1992; Tao, Cattaneo & Vainshtein 1993; Cattaneo 1994; and Cattaneo & Hughes 1996; see also Seehafer 1994, 1995) have argued that dynamo action is quenched if the large-scale magnetic field $\overline{B}$ in velocity units exceeds a critical value, $\overline{B}_c = R_M^{-1/2} v_0$, where $R_M = v_0 L / \lambda$ is the magnetic Reynolds number of the turbulence, v_0 is the turbulent velocity at the outer scale of the turbulence L , and $\lambda = \eta c^2 / 4\pi$ is the magnetic diffusivity, with η the resistivity. They argue that this result is supported by direct numerical simulations of MHD incompressible turbulence (Cattaneo & Hughes 1996). If they are correct, the classical α - Ω dynamo theory based upon Parker’s (1955) paper and developed by the Potsdam group (Krause and Rädler 1980; see also Moffatt 1978 and Parker 1979), which applies to weak large-scale magnetic fields, is not applicable to

present-day galaxies, as $\overline{B}$ is observed to greatly exceed $R_M^{-1/2}v_0$.

The argument by Cattaneo and his collaborators depends upon their analysis of nonlinear interactions which are surely present. The effects of such interactions have also been considered independently in a series of papers on MHD turbulence by Pouquet and her collaborators (Frisch, Pouquet, L  orat, & Mazure 1975; Pouquet, Frisch, & L  orat 1976; Pouquet & Patterson 1978; Meneguzzi, Frisch & Pouquet 1981) using spectral methods. By and large this work supports the applicability of the classical theory even for large-scale fields $\overline{B}$ approaching v_0 , which encompass those observed. Chandran (1996) confirms the results of Pouquet, Frisch, & L  orat (1976) using a different spectral method.

Since $R_M \gg 1$ in most astrophysical situations (as large as 10^{20} in the interstellar medium of our Galaxy) the difference between the critical field advocated by Cattaneo and his collaborators, $\overline{B}_c \sim R_M^{-1/2}v_0$, and that implied by Pouquet and her collaborators, $\overline{B}_c \sim v_0$, is crucial.

In this paper we extend the calculation of one of the dynamo coefficients, α , of the classical theory to include arbitrary values of $\overline{B}$. Our result contains two terms, one which resembles that advocated by Cattaneo and his collaborators in its dependence upon R_M . The other resembles a formula proposed by Kraichnan (1979) on the basis of a simple model incorporating damping by nonlinear interactions. For $R_M \gg 1$, the latter term dominates, so we find that the classical result for α applies for $\overline{B}$ of the order of v_0 . We plan to extend the simulations of Cattaneo and his collaborators to other values of R_M in order to clarify our disagreement with them.

We confine discussion to the alpha effect, and omit any discussion of turbulent diffusion, which is also controversial.

Before proceeding to the nonlinear theory we review the classical linear theory. The

classical theory is based on a clear separation of scales between the scale L of the dominant turbulent motions (here called the “outer scale”), and the size D of the system, with $L \ll D$. The large-scale magnetic field $\overline{\mathbf{B}}$ satisfies an induction equation,

$$\partial_t \overline{\mathbf{B}} = \nabla \times (\overline{\mathbf{V}} \times \overline{\mathbf{B}}) + \nabla \times \langle \mathbf{v} \times \mathbf{b} \rangle + \lambda \nabla^2 \overline{\mathbf{B}} , \quad (1)$$

where $\overline{\mathbf{V}}$ is the large-scale velocity field (differential rotation in the case of a galaxy), and $\langle \mathbf{v} \times \mathbf{b} \rangle$, called the “turbulent emf,” is the spatial average of the cross product of the small-scale velocity $\mathbf{v}$ and the small-scale magnetic field $\mathbf{b}$ over a scale much smaller than D , but much larger than L .

The first term in (1) leads to the so-called “ Ω -effect,” according to which lines of force of $\overline{\mathbf{B}}$ are stretched by the differential rotation, creating a growing toroidal field from a poloidal one. This term is not controversial, and will not be discussed further here. Controversy centers on the turbulent emf, $\langle \mathbf{v} \times \mathbf{b} \rangle$, which enables a growing toroidal field to feed back into the poloidal direction, giving exponential amplification of the large-scale $\overline{\mathbf{B}}$.

The evaluation of $\langle \mathbf{v} \times \mathbf{b} \rangle$ is usually restricted to incompressible isotropic turbulence. One might assume that if the turbulent velocity field $\mathbf{v}$ is isotropic, the small-scale magnetic field $\mathbf{b}$ would be isotropic also, so that the c -component of the emf would be

$$\langle \mathbf{v} \times \mathbf{b} \rangle_c = \epsilon_{cde} \langle v_d b_e \rangle = \frac{1}{3} \epsilon_{cde} \delta_{de} \langle \mathbf{v} \cdot \mathbf{b} \rangle = 0 , \quad (2)$$

in the light of an identity for isotropic tensors (Krause & Rädler 1980). However, this is not correct, because $\overline{\mathbf{B}}$, being anisotropic, induces a term in $\mathbf{b}$ that is not isotropic even though $\mathbf{v}$ is isotropic. Evaluating this term requires the induction equation for $\mathbf{b}$, which can be found by writing the induction equation for the total magnetic field $\mathbf{B} = \overline{\mathbf{B}} + \mathbf{b}$ and separating off the small-scale parts to give

$$\partial_t \mathbf{b} = \nabla \times (\mathbf{v} \times \overline{\mathbf{B}}) + \nabla \times (\overline{\mathbf{V}} \times \mathbf{b}) + \nabla \times (\mathbf{v} \times \mathbf{b}) - \nabla \times \langle \mathbf{v} \times \mathbf{b} \rangle + \lambda \nabla^2 \mathbf{b} . \quad (3)$$

The second term on the right in (3) represents a change of the reference frame to that moving at the mean velocity $\overline{\mathbf{V}}$, in which $\mathbf{v}$ may be assumed to be isotropic; we assume $\mathbf{V} = \text{const.}$, so we make that change and henceforth omit the term. The fourth term on the right is a large-scale quantity, so its time integral, being large scale, will contribute nothing when crossed with $\mathbf{v}$ (small-scale) and averaged, and so is neglected henceforth. The first term is $-\mathbf{v} \cdot \nabla \overline{\mathbf{B}} + \overline{\mathbf{B}} \cdot \nabla \mathbf{v}$. As the $\mathbf{v} \cdot \nabla \overline{\mathbf{B}}$ term ultimately leads to turbulent diffusion of $\overline{\mathbf{B}}$, which is not our main interest here, we omit it, leaving

$$\partial_t \mathbf{b} = \overline{\mathbf{B}} \cdot \nabla \mathbf{v} + \nabla \times (\mathbf{v} \times \mathbf{b}) + \lambda \nabla^2 \mathbf{b} . \quad (4)$$

The second term in (4) is neglected in the classical discussions in a step referred to as the first order smoothing approximation, or FOSA. Krause & Rädler (1980) showed that this is legitimate if λ is large, so R_M is small, because then Ohmic diffusion keeps b small, and the first term dominates the second. However, because R_M is large in astrophysics, this case is not relevant here. They also discuss the case when the Strouhal number

$$S = \frac{v_0 \tau}{L} \quad (5)$$

is small, where τ is the correlation time for eddies at the outer scale of the turbulence. Because $L/v_0 = t_{ed}$, the eddy turnover time, $S = \tau/t_{ed}$, and one might suppose that $S = O(1)$. In fact, it is observed experimentally that $S \simeq 0.2 - 0.3$ in ordinary hydrodynamic turbulence (Pope 1994). Although this value is not as small as one would like, it provides a way to approximate (4). To understand how, it is important to distinguish between $\mathbf{b}^{(0)}$, the small-scale magnetic field when $\overline{\mathbf{B}} = 0$, and $\mathbf{b}^{(1)}$, the perturbation to $\mathbf{b}$ when a small $\overline{\mathbf{B}}$ is present. If the turbulent velocity field $\mathbf{v}$ is isotropic, $\mathbf{b}^{(0)}$ is, and as shown in (2), its contribution to $\langle \mathbf{v} \times \mathbf{b} \rangle$ vanishes. However, $\mathbf{b}^{(1)}$ is not isotropic, and its contribution to $\langle \mathbf{v} \times \mathbf{b} \rangle$ does not vanish.

If we write $\mathbf{b} = \mathbf{b}^{(0)} + \mathbf{b}^{(1)}$, $\mathbf{b}^{(0)}$ is governed by

$$\partial_t \mathbf{b}^{(0)} = \nabla \times (\mathbf{v}^{(0)} \times \mathbf{b}^{(0)}) + \lambda \nabla^2 \mathbf{b}^{(0)}, \quad (6)$$

where $\mathbf{v}^{(0)}$ is the isotropic turbulent velocity. Parker (1979; p. 511) shows that if λ is small (the astrophysical case), (6) leads to exponential growth of $b^{(0)}$ with a time constant τ/S^2 until (in velocity units) it begins to approach $v^{(0)}$. At that point, we expect that back reaction due to the Lorentz force associated with the large value of the small-scale field $b^{(0)}$ will result in a steady state in which the energy driving the turbulence at the outer scale (buoyancy forces in stars, supernova explosions in galaxies) is balanced by a nonlinear cascade to smaller scales, where it is dissipated by viscosity and/or Joule heating. This expectation is confirmed by the Pouquet *et al.* (1976) and by direct simulations, *e.g.* Cattaneo and Vainshtein (1991), Cattaneo (1994), and Cattaneo and Hughes (1996). As we explain later, we will take this steady state as the base state which is perturbed by $\overline{\mathbf{B}}$. The first term in (6) mediates exchange of magnetic energy with kinetic energy in MHD turbulence, as discussed in Appendix A, and it certainly cannot be neglected.

However, we now argue that the corresponding term in the equation for $\mathbf{b}^{(1)}$ *can* be neglected, as follows. Evidently $\mathbf{b}^{(1)}$ is governed by the parts of (4) which are of first order in $\overline{\mathbf{B}}$, namely,

$$\partial_t \mathbf{b}^{(1)} = \overline{\mathbf{B}} \cdot \nabla \mathbf{v}^{(0)} + \nabla \times (\mathbf{v}^{(0)} \times \mathbf{b}^{(1)}) + \lambda \nabla^2 \mathbf{b}^{(1)}. \quad (7)$$

In the rest of this section, we assume that $R_M \gg 1$, so that we may neglect the third term. We formally integrate (7) to get

$$\begin{aligned} \mathbf{b}^{(1)}(\mathbf{x}, t) &= \overline{\mathbf{B}} \cdot \nabla \int_{-\infty}^t dt_1 \mathbf{v}^{(0)}(\mathbf{x}, t_1) \\ &\quad + \int_{-\infty}^t dt_1 \nabla \times [\mathbf{v}^{(0)}(\mathbf{x}, t_1) \times \mathbf{b}^{(1)}(\mathbf{x}, t_1)]. \end{aligned} \quad (8)$$

This does not seem useful, because the desired quantity, $\mathbf{b}^{(1)}$, appears under the integral. However, we can show that the integral is much smaller than $\mathbf{b}^{(1)}$ itself, which appears on the left-hand side, if $S \ll 1$.

Since $\mathbf{v}^{(0)}$ is a stochastic function of t , only those parts of the integral in (8) which come from times t_1 which differ from t by less than a correlation time τ will correlate significantly with $\mathbf{v}^{(0)}$ in the turbulent emf, $\langle \mathbf{v}^{(0)}(\mathbf{x}, t) \times \mathbf{b}^{(1)}(\mathbf{x}, t) \rangle$. Hence we can replace the lower limit on the integral by $t - \tau$, and then estimate the integral by τ times the integrand at $t_1 = t$. Since the order of magnitude of ∇ is L^{-1} , the order of magnitude of the second term in (8) is

$$\frac{\tau v^{(0)}}{L} b^{(1)}(t) = S b^{(1)}(t) , \quad (9)$$

whose ratio to the magnitude of the left hand side of (8) is S . If, as we shall argue later, $S < 1$, we can neglect the second term in (8), and hence in (7), so, still neglecting diffusion, (7) becomes

$$\partial_t \mathbf{b}^{(1)} = \overline{B} \cdot \nabla \mathbf{v}^{(0)} . \quad (10)$$

Effectively $b^{(1)} \sim S \overline{B}$, so if $S < 1$, $b^{(1)} < \overline{B}$, even when $\overline{B} \ll v^{(0)}$. However this says nothing about $b^{(0)}$, which as we have stressed, approaches $v^{(0)}$ in value, even if $v^{(0)} \gg \overline{B}$. Hopefully this discussion clarifies a point which has led to confusion in the past.

The solution to (10) can be written in component form as

$$b_e^{(1)}(t) = \overline{B}_p \partial_p \int_{-\infty}^t dt' v_e^{(0)}(t') , \quad (11)$$

so that

$$\langle \mathbf{v}^{(0)} \times \mathbf{b}^{(1)} \rangle_c = \epsilon_{cde} \left\langle v_d^{(0)}(t) \overline{B}_p \partial_p \int_{-\infty}^t dt' v_e^{(0)}(t') \right\rangle . \quad (12)$$

Averaging commutes with integration and differentiation, spatial differentiation commutes with time integration, and $\overline{B}$ is independent of time on the scale τ , so

$$\langle \nabla \times \mathbf{b} \rangle_c = \overline{B}_p \epsilon_{cde} \int_{-\infty}^t dt' \langle v_d^{(0)}(t) \partial_p v_e^{(0)}(t') \rangle . \quad (13)$$

Since $\mathbf{v}^{(0)}$ correlates with its derivatives for a time of order τ , the integral is of order τ times the average taken at a given time (Krause & Rädler 1980). To make further progress

in what follows, we need an explicit dependence of the indicated correlation on t and t' . Because the turbulence is assumed to be steady, it can depend only on $t - t'$, so a convenient representation is

$$\langle v_d^{(0)}(t) \partial_p v_e^{(0)}(t') \rangle = \langle v_d^{(0)} \partial_p v_e^{(0)} \rangle e^{-|t-t'|/\tau}, \quad (14)$$

where the common time argument in the second average has been omitted because the turbulence is presumed steady. Hence $\langle \mathbf{v} \times \mathbf{b} \rangle$ is given by

$$\langle \mathbf{v} \times \mathbf{b} \rangle_c = \overline{B}_p \tau \epsilon_{cde} \langle v_d^{(0)} \partial_p v_e^{(0)} \rangle. \quad (15)$$

At this point we use the assumption that the velocity $\mathbf{v}^{(0)}$ is distributed isotropically. According to Krause & Rädler (1980), a third-rank isotropic tensor like that in (15) can be written

$$\langle v_d^{(0)} \partial_p v_e^{(0)} \rangle = \frac{1}{6} \epsilon_{dpe} \langle \mathbf{v}^{(0)} \cdot \nabla \times \mathbf{v}^{(0)} \rangle, \quad (16)$$

so from (13),

$$\langle \mathbf{v} \times \mathbf{b} \rangle = \alpha \overline{\mathbf{B}}, \quad (17)$$

where

$$\alpha = -\frac{1}{3} \tau \langle \mathbf{v}^{(0)} \cdot \nabla \times \mathbf{v}^{(0)} \rangle \quad (18)$$

is the classical expression for the dynamo coefficient (Krause & Rädler 1980, Moffatt 1978), but with the additional feature that the turbulent velocities indicated refer to the zero-order state. The quantity in angular brackets, a pseudoscalar, is known as the kinetic helicity of the turbulent flow $\mathbf{v}^{(0)}$.

In an independent development, Pouquet, Frisch & L  orat (1976) calculated magnetic energy spectra for MHD turbulence, solving the spectral equations using a closure method known as the EDQNM (Eddy-Damped Quasi-Normal Markovian) approximation. They found that if $\langle \mathbf{v}^{(0)} \cdot \nabla \times \mathbf{v}^{(0)} \rangle$ vanishes, the magnetic energy spectrum E_k^M peaks near k_0 , the wave number at which turbulent energy is injected, and reaches a steady state in

which the total energy E_k , the sum of the kinetic energy E_k^V and the magnetic energy E_k^M , cascades to higher wave numbers, ultimately to be lost to Ohmic dissipation and/or viscosity.

If, on the other hand, $\langle \mathbf{v}^{(0)} \cdot \nabla \times \mathbf{v}^{(0)} \rangle \neq 0$, they found that E_k^M inverse cascades, accumulating at an ever-decreasing wave number. They attributed this to the turbulent dynamo effect described above, operating in the nonlinear regime. In the special case that the wave number k of interest is $\ll k_0$, so that there is a clear separation of scales, they find an approximate expression for a quantity α_k governing the growth of E_k^M , where

$$\alpha_k = -\frac{4}{3} \int_{k/a}^{\infty} dq \theta_{kqq} (H_q^V - q^2 H_q^M) . \quad (19)$$

Here a is a small parameter, θ_{kqq} is an effective correlation time for modes of wave number q , and H_q^V and $q^2 H_q^M$ are the spectra corresponding to the kinetic helicity correlation function $\langle \mathbf{v}^{(0)}(\mathbf{x}) \cdot \nabla \times \mathbf{v}^{(0)}(\mathbf{x} + \boldsymbol{\xi}) \rangle$ and the current (Keinigs 1983) helicity correlation function $\langle \mathbf{b}^{(0)}(\mathbf{x}) \cdot \nabla \times \mathbf{b}^{(0)}(\mathbf{x} + \boldsymbol{\xi}) \rangle$, respectively. (Here $\mathbf{b}$ is in velocity units, obtained by dividing $\mathbf{b}$ by $\sqrt{4\pi\rho}$, so that $\mathbf{b}$ is the vector Alfvén velocity.) The term adopted by Pouquet *et al.* for (19), the *residual torsality*, has not reappeared in the literature. Here we note that the first term in (19) is similar to (18), so that classical dynamo theory can be interpreted as an inverse cascade in helical turbulence. The second term does not appear in the classical result, but Montgomery and Chen (1984) verified it using calculations in real rather than k space. Pouquet *et al.* thus found that dynamo action takes place in the fully nonlinear regime, with no restriction as to the magnitude of $\mathbf{b}$ or $\overline{\mathbf{B}}$, and that the classical expression for α must be modified by the addition of the second term in (19).

2. The Method of This Paper

In this paper we extend the classical analysis of the α effect into the regime of large $\overline{B}$. We use the standard two-scale approximation in real space, and at several points we refer to the results obtained by Pouquet *et al.* (1976) using the spectral closure method. We find that when a clear distinction is made between the quantities $\mathbf{v}$ and $\mathbf{b}$ on the one hand, and their values when $\overline{\mathbf{B}} = 0$, $\mathbf{v}^{(0)}$ and $\mathbf{b}^{(0)}$, nonlinearity due to $\overline{B}$ appears in a straightforward manner. A key feature of our derivation is the assumption that correlations are damped by nonlinear interactions, following Pouquet *et al.* (1976) and Kraichnan (1979). As shown by Kichanitov (1985), such an assumption can be justified by application of renormalization group methods.

Our results depend upon the damping rate at each wave number k , γ_k , the spatial spectra of $\mathbf{v}^{(0)}$ and $\mathbf{b}^{(0)}$, the value of $\overline{B}$, and the value of λ , the magnetic diffusivity, expressed in terms of the magnetic Reynolds number R_M , which for convenience we assume to be equal to the Reynolds number R . We find that α is a well-behaved function of the parameters, and that the behavior for larger R_M is similar to that predicted by Kraichnan (1979).

We take as our base state fully-developed MHD turbulence driven by external forces and in a steady state as a result of a turbulent cascade to large wave numbers, but with $\overline{B} = 0$. As demonstrated by Pouquet *et al.* (1976), in such turbulence there is approximate equipartition between the magnetic energy E_k^M and the kinetic energy E_k^V for wave numbers $k \gtrsim 3k_0$, where $L = k_0^{-1}$ is the outer scale of the turbulence. As shown by Parker (1979, p. 513), approach to the steady state occurs on the scale of the eddy turnover time $t_{ed} = L/v^{(0)}$, which, as stressed by Kulsrud & Anderson (1992), is much shorter than the dynamo growth time. According to the calculations of Pouquet *et al.*, the saturation which occurs at small scales (large k) does not prevent the increase of magnetic energy on scales

larger than L if the turbulence is helical. It is important to note that the growth of a large-scale field as a consequence of an α effect does not substantially modify the spectra of E_k^M calculated by Pouquet *et al.* (1976) for k of the order of k_0 (their Fig. 8), so our concept of a base state independent of $\bar{\mathbf{B}}$ is valid. We assume that the properties of the base state can be calculated once and for all by numerical simulations. Our results then allow us to calculate α in terms of those properties.

Following Montgomery and Chen (1984), we present our calculations in terms of the Elsässer variables $\mathbf{z}^\pm = \mathbf{v} \pm \mathbf{b}$, (Biskamp 1993) and have checked our results by carrying out the calculation in terms of $\mathbf{v}$ and $\mathbf{b}$. Elsässer variables are naturally adapted to the problem, shortening the calculation substantially. More important, we show in Appendix A that unlike the kinetic energy E_k^V associated with $\mathbf{v}$ and the magnetic energy E_k^M associated with $\mathbf{b}$, both E_k^+ and E_k^- cascade directly in isotropic turbulence, allowing us to employ a single damping constant $\gamma_k = \gamma_k^+ = \gamma_k^-$ to describe the effects of the nonlinear terms in the base state.

Our goal is to calculate the turbulent emf, $\langle \mathbf{v} \times \mathbf{b} \rangle$. Since

$$\mathbf{v} = \frac{1}{2}(\mathbf{z}^+ + \mathbf{z}^-) \quad (20)$$

and

$$\mathbf{b} = \frac{1}{2}(\mathbf{z}^+ - \mathbf{z}^-) , \quad (21)$$

we have

$$\begin{aligned} \langle \mathbf{v} \times \mathbf{b} \rangle_c &= \frac{1}{4} \left\langle (\mathbf{z}^+ + \mathbf{z}^-) \times (\mathbf{z}^+ - \mathbf{z}^-) \right\rangle_c = -\frac{1}{2} \left\langle \mathbf{z}^+ \times \mathbf{z}^- \right\rangle_c \\ &= -\frac{1}{2} \left\langle \epsilon_{cde} z_d^+ z_e^- \right\rangle . \end{aligned} \quad (22)$$

We use this formula in what follows.

3. Evolution of the Turbulent Fields

As explained above, the correlation indicated in (22) vanishes for isotropic turbulence, but is nonzero when one takes into account the perturbations of $\mathbf{z}^+$ and $\mathbf{z}^-$ which are caused by $\overline{\mathbf{B}}$. To obtain these, we consider the dynamical equations for $\mathbf{z}^\pm$, or equivalently, $\mathbf{v}$ and $\mathbf{b}$.

When Ohmic dissipation is included, the induction equation for $\mathbf{b}$ is, from (3),

$$\partial_t \mathbf{b} = -\mathbf{v} \cdot \nabla \overline{\mathbf{B}} + \overline{\mathbf{B}} \cdot \nabla \mathbf{v} - \overline{\mathbf{V}} \cdot \nabla \mathbf{b} + \mathbf{b} \cdot \nabla \overline{\mathbf{V}} - \mathbf{v} \cdot \nabla \mathbf{b} + \mathbf{b} \cdot \nabla \mathbf{v} + \lambda \nabla^2 \mathbf{b} , \quad (23)$$

where λ is the magnetic diffusivity. As explained in §1, adopting in a frame of reference moving with $\overline{\mathbf{V}}$ eliminates the third term on the right, and the fourth term can be neglected with respect to the sixth because we assume that the size of the system $S \gg L$. The remaining terms can be written in terms of $\mathbf{B} = \overline{\mathbf{B}} + \mathbf{b}$ as

$$\partial_t \mathbf{b} = -\mathbf{v} \cdot \nabla \mathbf{B} + \mathbf{B} \cdot \nabla \mathbf{v} + \lambda \nabla^2 \mathbf{b} . \quad (24)$$

The classical theory ignores the effect of $\overline{\mathbf{B}}$ on the velocity, on the grounds that the Lorentz force associated with $\overline{\mathbf{B}}$ is of order $\overline{\mathbf{B}}^2$, hence negligible in the limit $\overline{\mathbf{B}} \rightarrow 0$. However, as we have explained above, $b^{(0)}$ grows quickly to approximate equipartition, so even in a first-order calculation, a Lorentz force proportional to $\overline{\mathbf{B}} b^{(0)}$ must be included. It is therefore essential to consider the effect of $\overline{\mathbf{B}}$ on $\mathbf{v}$; we shall do so to all orders in $\overline{\mathbf{B}}$. To do this, we use the equation of motion for the small-scale velocity $\mathbf{v}$:

$$\partial_t \mathbf{v} = -\mathbf{v} \cdot \nabla \mathbf{v} - \nabla p + \mathbf{B} \cdot \nabla \mathbf{B} - \nabla \frac{1}{2} B^2 + \nu \nabla^2 \mathbf{v} + \mathbf{f} , \quad (25)$$

where $\mathbf{f}$ is the applied force per unit mass and ν is the kinematic viscosity; $\mathbf{B}$, the magnetic field $\div \sqrt{4\pi\rho}$, is in velocity units. Hence

$$\partial_t \mathbf{v} = -\mathbf{v} \cdot \nabla \mathbf{v} - \nabla P + \mathbf{B} \cdot \nabla \mathbf{B} + \nu \nabla^2 \mathbf{v} + \mathbf{f} , \quad (26)$$

where

$$P = p + \frac{1}{2}B^2 . \quad (27)$$

We define the Elssäser variables for the field $\mathbf{B}$ as

$$\mathbf{Z}^\pm = \mathbf{v} \pm \mathbf{B} = \mathbf{v} \pm \mathbf{b} \pm \overline{\mathbf{B}} = \mathbf{z}^\pm \pm \overline{\mathbf{B}} , \quad (28)$$

so that

$$\mathbf{v} = \frac{1}{2}(\mathbf{Z}^+ + \mathbf{Z}^-) \quad (29)$$

and

$$\mathbf{B} = \mathbf{b} + \overline{\mathbf{B}} = \frac{1}{2}(\mathbf{Z}^+ - \mathbf{Z}^-) . \quad (30)$$

Then (24) becomes

$$\begin{aligned} \frac{1}{2}\partial_t \mathbf{Z}^+ - \frac{1}{2}\partial_t \mathbf{Z}^- &= -\frac{1}{4}(\mathbf{Z}^+ + \mathbf{Z}^-) \cdot \nabla(\mathbf{Z}^+ - \mathbf{Z}^-) \\ &\quad + \frac{1}{4}(\mathbf{Z}^+ - \mathbf{Z}^-) \cdot \nabla(\mathbf{Z}^+ + \mathbf{Z}^-) \\ &\quad + \frac{1}{2}\lambda \nabla^2(\mathbf{Z}^+ - \mathbf{Z}^-) . \end{aligned} \quad (31)$$

(Note that we have set $\partial_t \overline{\mathbf{B}} = 0$ because $\overline{\mathbf{B}}$ varies only on the long time scale.) Equation (26) becomes

$$\begin{aligned} \frac{1}{2}\partial_t \mathbf{Z}^+ + \frac{1}{2}\partial_t \mathbf{Z}^- &= -\frac{1}{4}(\mathbf{Z}^+ + \mathbf{Z}^-) \cdot \nabla(\mathbf{Z}^+ + \mathbf{Z}^-) \\ &\quad + \frac{1}{4}(\mathbf{Z}^+ - \mathbf{Z}^-) \cdot \nabla(\mathbf{Z}^+ - \mathbf{Z}^-) \\ &\quad + \frac{1}{2}\nu \nabla^2(\mathbf{Z}^+ + \mathbf{Z}^-) - \nabla P + \mathbf{f} . \end{aligned} \quad (32)$$

Adding and subtracting (31) and (32) yields

$$\partial_t \mathbf{Z}^\pm = -\mathbf{Z}^\mp \cdot \nabla \mathbf{Z}^\pm + \frac{1}{2}(\nu + \lambda) \nabla^2 \mathbf{Z}^\pm + \frac{1}{2}(\nu - \lambda) \nabla^2 \mathbf{Z}^\mp - \nabla P + \mathbf{f} . \quad (33)$$

The dissipative coupling between $\mathbf{Z}^+$ and $\mathbf{Z}^-$ vanishes in the special case that the magnetic Prandtl number λ/ν equals unity. For simplicity we assume that is the case in what follows;

we note that both Pouquet *et al.* (1976) and Cattaneo and Hughes (1996) also made the same assumption. Chou and Fish (1998) discuss the case $\lambda/\nu \neq 1$.

From (28) the nonlinear term in (33) is

$$\begin{aligned} \mathbf{z}^\mp \cdot \nabla \mathbf{z}^\pm &= (\mathbf{z}^\mp \mp \overline{\mathbf{B}}) \cdot \nabla (\mathbf{z}^\pm \pm \overline{\mathbf{B}}) \\ &= \mathbf{z}^\mp \cdot \nabla \mathbf{z}^\pm \mp \overline{\mathbf{B}} \cdot \nabla \mathbf{z}^\pm, \end{aligned} \quad (34)$$

where we have neglected $\nabla \overline{\mathbf{B}}$ for the reasons given previously. Hence (33) becomes

$$\partial_t \mathbf{z}^\pm = -\mathbf{z}^\mp \cdot \nabla \mathbf{z}^\pm + \lambda \nabla^2 \mathbf{z}^\pm - \nabla P + \mathbf{f} \pm \overline{\mathbf{B}} \cdot \nabla \mathbf{z}^\pm, \quad (35)$$

where we have neglected $\partial_t \overline{\mathbf{B}}$ for the reason given previously.

We adopt a perturbation expansion in $\overline{\mathbf{B}}$, in which the zero-order variables $\mathbf{z}^{\pm(0)}$ describe the turbulence exactly if $\overline{\mathbf{B}}$ is zero. Therefore $\mathbf{z}^{\pm(0)}$ satisfies

$$\partial_t \mathbf{z}^{\pm(0)} = -\mathbf{z}^{\mp(0)} \cdot \nabla \mathbf{z}^{\pm(0)} + \lambda \nabla^2 \mathbf{z}^{\pm(0)} - \nabla P^{(0)} + \mathbf{f}, \quad (36)$$

where

$$P^{(0)} = p^{(0)} + \frac{1}{2} B^2|_{\overline{\mathbf{B}}=0} = p^{(0)} + \frac{1}{2} (b^{(0)})^2. \quad (37)$$

If we assume that the driving force is independent of $\overline{\mathbf{B}}$, $\mathbf{f}$ has the same value in both the zero-order and the perturbed state.

Equation (36) can be solved numerically for a variety of initial conditions, and so in principle any averages of zero-order quantities required can be computed. Our task is then to compute $\langle \mathbf{v} \times \mathbf{b} \rangle$, which depends on the perturbations induced by $\overline{\mathbf{B}}$, in terms of averages over zero-order quantities.

We let

$$\mathbf{z}^\pm = \mathbf{z}^{\pm(0)} + \mathbf{z}'^\pm, \quad (38)$$

where $\mathbf{z}'^{\pm}$ contains perturbations of all orders in $\overline{B}$. Although in principle $\mathbf{z}'^{\pm}$ can be represented as a power series in $\overline{B}$, we find that it is not necessary to do so for the special case $\lambda/\nu = 1$. If we eliminate $\mathbf{f}$ between (35) and (36), we find that

$$\begin{aligned} (\partial_t - \lambda \nabla^2) \mathbf{z}'^{\pm} &= -\mathbf{z}'^{\mp} \cdot \nabla \mathbf{z}^{\pm(0)} - \mathbf{z}^{\mp(0)} \cdot \nabla \mathbf{z}'^{\pm} - \mathbf{z}'^{\mp} \cdot \nabla \mathbf{z}'^{\pm} \\ &\quad - \nabla(P - P^{(0)}) \pm \overline{\mathbf{B}} \cdot \nabla \mathbf{z}^{\pm} . \end{aligned} \quad (39)$$

Inspecting (39), we see that the left hand side and the term $\overline{\mathbf{B}} \cdot \nabla \mathbf{z}'^{\pm}$ on the right are linear in $\mathbf{z}'^{\pm}$, hence easy to deal with. If R_M is moderately large, we can ignore the term $\lambda \nabla^2 \mathbf{z}'^{\pm}$ in assessing the order of magnitude of the remaining terms. Putting aside the pressure term for the moment, we can write (39) in the symbolic form

$$\partial_t z' = L^{-1} \left[z' z^{(0)}, (z')^2, \overline{B} z^{(0)}, \overline{B} z' \right] , \quad (40)$$

where the commas separate terms of potentially different orders, and where we have combined the first two terms on the right of (39) into one. As in our discussion of the classical case, we take $\Delta z' = \int dt \partial_z z'$, the change in z' after one correlation time τ , to equal z' in order of magnitude, by definition. Then (40) implies that

$$z' = \frac{\tau}{t_{ed}} \left[z', (z')^2/z^{(0)}, \overline{B}, \overline{B} z'/z^{(0)} \right] , \quad (41)$$

where $t_{ed} \equiv L/z^{(0)} \simeq L/v^{(0)}$.

In the classical discussion it is assumed that because $\overline{B}$ is small, back reaction by the Lorentz force can be neglected, so the motions are hydrodynamic in nature. That allowed us to use the fact that τ/t_{ed} is small (see below). That is not really true, because, as we discussed earlier, the zero-order state quickly approaches equipartition. We want to argue, however, that even in the nonlinear case τ/t_{ed} is a small number. To do so, we appeal to the nonlinear calculations of Pouquet *et al.* (1976) regarding the zero-order state. They showed that the magnetic energy spectrum E_k^M is in equipartition with the kinetic energy

spectrum E_k^V for $k \gtrsim 3k_0$, where k_0 is the wave number at which kinetic energy is being injected (the outer scale of the turbulence). In this range, back reaction is a major effect, causing the turbulence to become a field of interacting Alfvén waves. However, these large wave numbers need not concern us, because in such waves, the current is perpendicular to the small-scale field, and the vorticity is perpendicular to the small-scale velocity, so that the two helicities vanish.

Of crucial importance for us, Pouquet *et al.* (1976) found that in the range $k_0 \lesssim k \lesssim 3k_0$, $E_k^M < E_k^V$, so that the motions are largely hydrodynamic in character, with only a modest back reaction of the magnetic field. This range is important because it contains most of the energy. Pope (1994) states that in a pure hydrodynamic turbulence it is experimentally observed that

$$\frac{\tau}{t_{ed}} = 0.2 - 0.3 . \quad (42)$$

We assume that (42) applies to the energy-containing eddies in MHD turbulence driven like that of Pouquet *et al.* Although τ/t_{ed} is not a very small number, we may take it to be a small parameter ϵ in analyzing (41). Thus, (41) becomes

$$z' \sim \epsilon \left[z', (z')^2/z^{(0)}, \overline{B}, \overline{B}z'/z^{(0)} \right] . \quad (43)$$

It seems reasonable to assume that the balance in (43) is between z' and the third term on the right, so

$$z' \sim \epsilon \overline{B} . \quad (44)$$

We check this by evaluating the first two terms under that assumption. We see that the ratio of the first to the third terms, and the ratio of the second to the fourth term, are both ϵ , so the first and second terms can be neglected as a first approximation. Note, however, that the ratio of the fourth to the third term is

$$z'/z^{(0)} = \epsilon \overline{B}/z^{(0)} \sim \epsilon \overline{B}/v^{(0)} , \quad (45)$$

which cannot be neglected because, although ϵ is small, we want a result which is valid to all orders in $\overline{B}/v^{(0)}$. Indeed, this is the source of the nonlinearity in our calculation.

Note that although we have used the short correlation time approximation to simplify our equations for z' , no assumption is made regarding the ratio of b to $\overline{B}$. This is a step forward, because the classical discussion has been justly criticized for ignoring back reaction, which amounts to assuming that b is small.

We now apply the divergence operator $\nabla \cdot$ to (39) with the first three terms on the right dropped. Because $\overline{\mathbf{B}}$ is constant and $\nabla \cdot$ commutes with $\overline{\mathbf{B}} \cdot \nabla$, the fact that

$$\nabla \cdot \mathbf{z}^\pm = \nabla \cdot (\mathbf{v} \pm \mathbf{b}) = 0 \quad (46)$$

for incompressible turbulence then implies that

$$\nabla^2(P - P^{(0)}) = 0 , \quad (47)$$

which for a homogeneous system implies that $P - P^{(0)} = \text{const.}$, so that

$$\nabla(P - P^{(0)}) = 0 . \quad (48)$$

Hence (39) becomes

$$(\partial_t - \lambda \nabla^2 \mp \overline{\mathbf{B}} \cdot \nabla) \mathbf{z}'^\pm = \pm \overline{\mathbf{B}} \cdot \nabla \mathbf{z}^{\pm(0)} . \quad (49)$$

We must solve this equation in order to calculate the turbulent emf according to (22).

The operator $\overline{\mathbf{B}} \cdot \nabla$ is best handled by introducing the spatial Fourier transform

$$\hat{\mathbf{z}}'^\pm(\mathbf{k}, t) = (2\pi)^{-3} \int d\mathbf{x} e^{-i\mathbf{k} \cdot \mathbf{x}} \mathbf{z}'^\pm(\mathbf{x}, t) , \quad (50)$$

so that when written in terms of the components of $\mathbf{z}$, (49) becomes

$$(\partial_t + \lambda k^2 \mp i\mathbf{k} \cdot \overline{\mathbf{B}}) \hat{z}'^\pm_d(\mathbf{k}, t) = \pm i\mathbf{k} \cdot \overline{\mathbf{B}} \hat{z}^{\pm(0)}_d(\mathbf{k}, t) , \quad (51)$$

to which the solution is

$$\hat{z}_d^{\pm}(\mathbf{k}, t) = \pm i \mathbf{k} \cdot \bar{\mathbf{B}} \int_{-\infty}^t dt_1 \exp \left\{ - \left[\mp i \mathbf{k} \cdot \bar{\mathbf{B}} + \lambda k^2 \right] (t - t_1) \right\} \hat{z}_d^{\pm(0)}(\mathbf{k}, t_1) , \quad (52)$$

where we have put the lower limit equal to $-\infty$ because we will find that the short correlation time makes values of t_1 significantly smaller than t irrelevant. The nonlinear dependence on $\bar{\mathbf{B}}$ is evident here.

From (22) and (38), the term of zero order in $z^{\pm}$ drops out according to (2), leaving

$$\begin{aligned} \langle \mathbf{v} \times \mathbf{b} \rangle_c &= -\frac{1}{2} \left\langle \epsilon_{cde} \hat{z}_d^{+(0)}(\mathbf{x}, t) \hat{z}_e'^-(\mathbf{x}, t) \right\rangle - \frac{1}{2} \left\langle \epsilon_{cde} \hat{z}_d'^+(\mathbf{x}, t) \hat{z}_e^{-(0)}(\mathbf{x}, t) \right\rangle \\ &\quad - \frac{1}{2} \left\langle \epsilon_{cde} \hat{z}_d'^+(\mathbf{x}, t) \hat{z}_e'^-(\mathbf{x}, t) \right\rangle . \end{aligned} \quad (53)$$

Inverting the Fourier transforms, we see from (50) that

$$\hat{z}_d^{\pm(0)}(\mathbf{x}, t) = \int d\mathbf{k} e^{i\mathbf{k} \cdot \mathbf{x}} \hat{z}_d^{\pm(0)}(\mathbf{k}, t) , \quad (54)$$

and, from (52)

$$\hat{z}_e'^{\pm}(\mathbf{x}, t) = \pm i \int d\mathbf{k}' e^{i\mathbf{k}' \cdot \mathbf{x}} \mathbf{k}' \cdot \bar{\mathbf{B}} \int_{-\infty}^t dt_1 \exp \left\{ - \left[\mp i \mathbf{k}' \cdot \bar{\mathbf{B}} + \lambda k'^2 \right] (t - t_1) \right\} \hat{z}_e^{\pm(0)}(\mathbf{k}', t_1) . \quad (55)$$

Hence the first term in (53) is

$$\frac{1}{2} i \epsilon_{cde} \left\langle \int d\mathbf{k} \int d\mathbf{k}' e^{i(\mathbf{k} + \mathbf{k}') \cdot \mathbf{x}} \mathbf{k}' \cdot \bar{\mathbf{B}} \int_{-\infty}^t dt_1 \exp \left\{ - [i \mathbf{k}' \cdot \bar{\mathbf{B}} + \lambda k'^2] (t - t_1) \right\} \hat{z}_d^{+(0)}(\mathbf{k}, t) \hat{z}_e^{-(0)}(\mathbf{k}', t_1) \right\rangle . \quad (56)$$

Because averaging commutes with integration, this can be written

$$\frac{1}{2} i \int d\mathbf{k} \int d\mathbf{k}' e^{i(\mathbf{k} + \mathbf{k}') \cdot \mathbf{x}} \mathbf{k}' \cdot \bar{\mathbf{B}} \int_{-\infty}^t dt_1 \exp \left\{ - [i \mathbf{k}' \cdot \bar{\mathbf{B}} + \lambda k'^2] (t - t_1) \right\} \left\langle \epsilon_{cde} \hat{z}_d^{+(0)}(\mathbf{k}, t) \hat{z}_e^{-(0)}(\mathbf{k}', t_1) \right\rangle . \quad (57)$$

The indicated correlation decreases rapidly to zero as $\mathbf{k} - \mathbf{k}'$ and $t - t_1$ go to zero. In Appendix B we show that the following representation is a reasonable one:

$$\left\langle \epsilon_{cde} \hat{z}_d^{+(0)}(\mathbf{k}, t) \hat{z}_e^{-(0)}(\mathbf{k}', t_1) \right\rangle = \epsilon_{cde} R_{de}(-\mathbf{k}) \delta(\mathbf{k} + \mathbf{k}') e^{-\gamma_k |t - t_1|} . \quad (58)$$

In principle, γ_k could be different for $\mathbf{z}^+$ and $\mathbf{z}^-$ modes, but we show in Appendix A that in turbulence in which the cross helicity $K = \frac{1}{2} \langle \mathbf{v} \cdot \mathbf{B} \rangle$, an ideal invariant, vanishes, the associated energies E^+ and E^- cascade directly and are equal. It is therefore reasonable to assume that if $K = 0$, $\gamma_k^+ = \gamma_k^-$. Following Pouquet *et al.* we assume that $K = 0$, and thus, that $\gamma_k^+ = \gamma_k^-$. Note that because γ_k is a parameter of the zero-order state, it does not depend upon $\overline{B}$. From Krause and Rädler (1980, p. 75) an isotropic tensor like $R_{de}(\mathbf{k})$ can be expressed in the form

$$\begin{aligned} R_{de}(\mathbf{k}) &= (A - k^{-1} \partial_k B) \delta_{de} - (\partial_k^2 B - k^{-1} \partial_k B) k^{-2} k_d k_e \\ &\quad + i k^{-1} \partial_k C \epsilon_{def} k_f, \end{aligned} \quad (59)$$

where A, B , and C are functions of k alone. Therefore

$$\epsilon_{cde} R_{de}(-\mathbf{k}) = -2i k_c k^{-1} \partial_k C \quad (60)$$

is nonvanishing only for turbulence lacking reflection symmetry, and so (58) becomes

$$\left\langle \epsilon_{cde} \hat{z}_d^{+(0)}(\mathbf{k}, t) \hat{z}_e^{-(0)}(\mathbf{k}', t_1) \right\rangle = -2i k_c k^{-1} \partial_k C e^{-\gamma_k |t - t_1|}. \quad (61)$$

The integration of (57) over $\mathbf{k}'$ converts $\mathbf{k}'$ to $-\mathbf{k}$, so $(\mathbf{k} + \mathbf{k}') \cdot \mathbf{x} = 0$. Since $|t - t_1| = t - t_1$ in the integration over t_1 , we find that (57) equals

$$\begin{aligned} & - \int d\mathbf{k} \mathbf{k} \cdot \overline{\mathbf{B}} k_c k^{-1} \partial_k C \int_{-\infty}^t dt_1 \exp \left\{ [i\mathbf{k} \cdot \overline{\mathbf{B}} - \lambda k^2 - \gamma_k](t - t_1) \right\} \\ & = i \int d\mathbf{k} \frac{i\mathbf{k} \cdot \overline{\mathbf{B}}}{(\lambda k^2 + \gamma_k) - i\mathbf{k} \cdot \overline{\mathbf{B}}} k_c k^{-1} \partial_k C. \end{aligned} \quad (62)$$

We note that the second term in (53) can be written

$$+ \frac{1}{2} \epsilon_{cde} \left\langle z_e'^+(\mathbf{x}, t) z_d^{-(0)}(\mathbf{x}, t) \right\rangle \quad (63)$$

because of the ϵ_{cde} . Then reference to (54) and (55) shows that this is the negative of (62), with $i \leftrightarrow -i$. Hence the sum of the first two terms in (53) can be written

$$i \int d\mathbf{k} k_c k^{-1} \partial_k C \left[\frac{i\mathbf{k} \cdot \overline{\mathbf{B}}}{(\lambda k^2 + \gamma_k) - i\mathbf{k} \cdot \overline{\mathbf{B}}} + \frac{i\mathbf{k} \cdot \overline{\mathbf{B}}}{(\lambda k^2 + \gamma_k) + i\mathbf{k} \cdot \overline{\mathbf{B}}} \right]$$

$$= -2 \int d\mathbf{k} k_c k^{-1} \partial_k C \mathbf{k} \cdot \bar{\mathbf{B}} (\lambda k^2 + \gamma_k) [(\lambda k^2 + \gamma_k)^2 + (\mathbf{k} \cdot \bar{\mathbf{B}})^2]^{-1} . \quad (64)$$

We now turn to the third term in (53), which is

$$\begin{aligned} & -\frac{1}{2} \int d\mathbf{k} \int d\mathbf{k}' e^{i(\mathbf{k}+\mathbf{k}') \cdot \mathbf{x}} (\mathbf{k} \cdot \bar{\mathbf{B}}) (\mathbf{k}' \cdot \bar{\mathbf{B}}) \\ & \cdot \int_{-\infty}^t dt_1 \exp \left\{ -[-i\mathbf{k} \cdot \bar{\mathbf{B}} + \lambda k^2](t - t_1) \right\} \\ & \cdot \int_{-\infty}^t dt_2 \exp \left\{ -[i\mathbf{k}' \cdot \bar{\mathbf{B}} + \lambda k'^2](t - t_2) \right\} \left\langle \epsilon_{cde} \hat{z}_d^{+(0)}(\mathbf{k}, t_1) \hat{z}_e^{-(0)}(\mathbf{k}', t_2) \right\rangle . \end{aligned} \quad (65)$$

According to (61), the average is

$$\left\langle \epsilon_{cde} \hat{z}_d^{+(0)}(\mathbf{k}, t_1) \hat{z}_e^{-(0)}(\mathbf{k}', t_2) \right\rangle = k_c k^{-1} \partial_k C \delta(\mathbf{k} + \mathbf{k}') e^{-\gamma_k |t_1 - t_2|} , \quad (66)$$

so integrating (65) over $\mathbf{k}'$ gives

$$\begin{aligned} & -i \int d\mathbf{k} (\mathbf{k} \cdot \bar{\mathbf{B}})^2 k_c k^{-1} \partial_k C \int_{-\infty}^t dt_1 \exp \left\{ -[-i\mathbf{k} \cdot \bar{\mathbf{B}} + \lambda k^2](t - t_1) \right\} \\ & \cdot \int_{-\infty}^t dt_2 \exp \left\{ -[-i\mathbf{k} \cdot \bar{\mathbf{B}} + \lambda k^2](t - t_2) \right\} e^{-\gamma_k |t_1 - t_2|} . \end{aligned} \quad (67)$$

In this expression, the integral over t_2 extends from $-\infty$ to t_1 with $|t_1 - t_2| = t_1 - t_2$, and from t_1 to t with $|t_1 - t_2| = t_2 - t_1$. We thus find that

$$\begin{aligned} & \int_{-\infty}^t dt_2 \exp \left\{ -[-i\mathbf{k} \cdot \bar{\mathbf{B}} + \lambda k^2](t_1 - t_2) - \gamma_k |t_1 - t_2| \right\} \\ & = \frac{2\gamma_k}{\gamma_k^2 - (-i\mathbf{k} \cdot \bar{\mathbf{B}} + \lambda k^2)^2} e^{i(\mathbf{k} \cdot \bar{\mathbf{B}} - \lambda k^2)(t - t_1)} + \frac{1}{-i\mathbf{k} \cdot \bar{\mathbf{B}} + \lambda k^2 - \gamma_k} e^{-\gamma_k(t - t_1)} , \end{aligned} \quad (68)$$

so that

$$\int_{-\infty}^t dt_1 (\quad) = \left(\frac{1}{i\mathbf{k} \cdot \bar{\mathbf{B}} - \lambda k^2} \right) \left(\frac{1}{i\mathbf{k} \cdot \bar{\mathbf{B}} - \lambda k^2 - \gamma_k} \right) , \quad (69)$$

where () refers to the function following dt_1 in (67). This expression can be written in the equivalent form

$$\int_{-\infty}^t dt_1 (\quad) = \gamma_k^{-1} \left[\frac{i\mathbf{k} \cdot \bar{\mathbf{B}} + \lambda k^2}{(\mathbf{k} \cdot \bar{\mathbf{B}})^2 + (\lambda k^2)^2} - \frac{i\mathbf{k} \cdot \bar{\mathbf{B}} + (\lambda k^2 + \gamma_k)}{(\mathbf{k} \cdot \bar{\mathbf{B}})^2 + (\lambda k^2 + \gamma_k)^2} \right] , \quad (70)$$

so (67) becomes

$$-i \int d\mathbf{k} (\mathbf{k} \cdot \overline{\mathbf{B}})^2 k_c k^{-1} \partial_k C \gamma_k^{-1} \left[\frac{i\mathbf{k} \cdot \overline{\mathbf{B}} + \lambda k^2}{(\lambda k^2)^2 + (\mathbf{k} \cdot \overline{\mathbf{B}})^2} - \frac{i\mathbf{k} \cdot \overline{\mathbf{B}} + (\lambda k^2 + \gamma_k)}{(\lambda k^2 + \gamma_k)^2 + (\mathbf{k} \cdot \overline{\mathbf{B}})^2} \right]. \quad (71)$$

We show below that the contributions of the terms of order $\overline{B}^2$ in the numerator vanish, so (71) becomes

$$\int d\mathbf{k} (\mathbf{k} \cdot \overline{\mathbf{B}})^3 k_c k^{-1} \partial_k C \gamma_k^{-1} \left[\frac{1}{(\lambda k^2)^2 + (\mathbf{k} \cdot \overline{\mathbf{B}})^2} - \frac{1}{(\lambda k^2 + \gamma_k)^2 + (\mathbf{k} \cdot \overline{\mathbf{B}})^2} \right]. \quad (72)$$

Combining (72) and (64) yields the following expression for (53):

$$\begin{aligned} \langle \mathbf{v} \times \mathbf{b} \rangle_c &= -2 \int d\mathbf{k} k_c k^{-1} \partial_k C \left\{ \frac{\mathbf{k} \cdot \overline{\mathbf{B}} (\lambda k^2 + \gamma_k)}{(\lambda k^2 + \gamma_k)^2 + (\mathbf{k} \cdot \overline{\mathbf{B}})^2} \right. \\ &\quad \left. - \frac{1}{2} (\mathbf{k} \cdot \overline{\mathbf{B}})^3 \gamma_k^{-1} \left[\frac{1}{(\lambda k^2)^2 + (\mathbf{k} \cdot \overline{\mathbf{B}})^2} - \frac{1}{(\lambda k^2 + \gamma_k)^2 + (\mathbf{k} \cdot \overline{\mathbf{B}})^2} \right] \right\}. \end{aligned} \quad (73)$$

To proceed, we adopt a coordinate system (1, 2, 3) such that

$$\overline{\mathbf{B}} = (\overline{B}, 0, 0) \quad (74)$$

has only one component, $\overline{B}_1 = \overline{B}$, so that

$$\mathbf{k} \cdot \overline{\mathbf{B}} = k_1 \overline{B}. \quad (75)$$

Hence

$$\begin{aligned} \langle \mathbf{v} \times \mathbf{b} \rangle_c &= -2 \int d\mathbf{k} k_c k^{-1} \partial_k C \left\{ \frac{k_1 \overline{B} (\lambda k^2 + \gamma_k)}{(\lambda k^2 + \gamma_k)^2 + (k_1 \overline{B})^2} \right. \\ &\quad \left. - \frac{1}{2} (k_1 \overline{B})^3 \gamma_k^{-1} \left[\frac{1}{(\lambda k^2)^2 + (k_1 \overline{B})^2} - \frac{1}{(\lambda k^2 + \gamma_k)^2 + (k_1 \overline{B})^2} \right] \right\}. \end{aligned} \quad (76)$$

Note that $\langle \mathbf{v} \times \mathbf{b} \rangle$ vanishes if $\overline{B} = 0$; as explained earlier, finite $\overline{B}$ is required to break rotational symmetry.

In our coordinate system

$$d\mathbf{k} = k^2 \sin \theta d\varphi d\theta dk , \quad (77)$$

where $\theta = \angle_{(1)}^{\mathbf{k}}$, and φ is the azimuth in the 2-3 plane. If $c = 2$ or 3 in (76), $k_c = k \sin \theta \sin \varphi$ or $k \sin \theta \cos \varphi$, and as the rest of the integrand is independent of φ , the integral over φ vanishes. We are left with $c = 1$, in which case the integrand is an even function of k_1 . The integral over φ gives 2π , so

$$d\mathbf{k} \rightarrow -2\pi k^2 dk d(\cos \theta) = -2\pi k dk dk_1 , \quad (78)$$

where we have changed variables from (k, θ) to $(k, k \cos \theta) = (k, k_1)$. As the integral over k_1 is from k to $-k$, and the integrand is even in k_1 ,

$$d\mathbf{k} \rightarrow 4\pi k dk dk_1 , \quad (79)$$

with k_1 varying from 0 to k . As claimed previously, the terms of order $\overline{B}^2$ in (71) would have contributed terms of order k_1^3 to the integrand in (72); as this is odd in k_1 , their contribution would have vanished when integrated from $-k$ to k . Hence

$$\begin{aligned} \langle \mathbf{v} \times \mathbf{b} \rangle_1 &= -2 \int_{k=0}^{\infty} 4\pi k dk \int_{k_1=0}^k dk_1 k_1 k^{-1} \partial_k C \left\{ \frac{k_1 \overline{B} (\lambda k^2 + \gamma_k)}{(\lambda k^2 + \gamma_k)^2 + (k_1 \overline{B})^2} \right. \\ &\quad \left. - \frac{1}{2} (k_1 \overline{B})^3 \gamma_k^{-1} \left[\frac{1}{(\lambda k^2)^2 + (k_1 \overline{B})^2} - \frac{1}{(\lambda k^2 + \gamma_k)^2 + (k_1 \overline{B})^2} \right] \right\} . \end{aligned} \quad (80)$$

Hence $\langle \mathbf{v} \times \mathbf{b} \rangle$ is parallel to $\overline{\mathbf{B}}$ as in the result of the classical theory, (17), and the dynamo coefficient α is

$$\begin{aligned} \alpha &= -8\pi \int_{k=0}^{\infty} dk \partial_k C \int_{k_1=0}^k dk_1 \left\{ \frac{k_1^2 (\lambda k^2 + \gamma_k)}{(\lambda k^2 + \gamma_k)^2 + (k_1 \overline{B})^2} \right. \\ &\quad \left. - \frac{1}{2} k_1^4 \overline{B}^2 \gamma_k^{-1} \left[\frac{1}{(\lambda k^2)^2 + (k_1 \overline{B})^2} - \frac{1}{(\lambda k^2 + \gamma_k)^2 + (k_1 \overline{B})^2} \right] \right\} . \end{aligned} \quad (81)$$

To carry out the integration over k_1 in (81), in the first term in braces we set

$$\xi = \frac{k \overline{B}}{\lambda k^2 + \gamma_k} , \quad (82)$$

so that

$$(\lambda k^2 + \gamma_k) \int_{k_1=0}^k dk_1 \frac{k_1^2}{(\lambda k^2 + \gamma_k)^2 + (k_1 \bar{B})^2} = k^3 (\lambda k^2 + \gamma_k)^{-1} \frac{\xi - \tan^{-1} \xi}{\xi^3} . \quad (83)$$

In the first term in brackets we set

$$\eta = \frac{k \bar{B}}{\lambda k^2} , \quad (84)$$

so that

$$-\frac{1}{2} \bar{B}^2 \gamma_k^{-1} \int_0^k \frac{k_1^4 dk_1}{(\lambda k^2 + \gamma_k)^2 + (k_1 \bar{B})^2} = -\frac{1}{2} \gamma_k^{-1} k^3 \left(\frac{\frac{1}{3} \eta^3 - \eta + \tan^{-1} \eta}{\eta^3} \right) . \quad (85)$$

In the second term in brackets, we again use (82), so that

$$\frac{1}{2} \bar{B}^2 \gamma_k^{-1} \int_0^k \frac{k_1^4 dk_1}{(\lambda k^2 + \gamma_k)^2 + (k_1 \bar{B})^2} = \frac{1}{2} \gamma_k^{-1} k^3 \left(\frac{\frac{1}{3} \xi^3 - \xi + \tan^{-1} \xi}{\xi^3} \right) . \quad (86)$$

Combining (83), (85), and (86), we find that in (81)

$$\int_{k_1=0}^k dk_1 \{ \} = \frac{1}{2} k^3 \gamma_k^{-1} \left[\frac{\gamma_k - \lambda k^2}{\gamma_k + \lambda k^2} f(\xi) + f(\eta) \right] , \quad (87)$$

where

$$f(\xi) = \frac{\xi - \tan^{-1} \xi}{\xi^3} , \quad (88)$$

so that (81) becomes

$$\alpha = - \int_{k=0}^{\infty} 4\pi k^2 dk k \partial_k C \gamma_k^{-1} \left[\frac{\gamma_k - \lambda k^2}{\gamma_k + \lambda k^2} f(\xi) + f(\eta) \right] . \quad (89)$$

Since the integrand is a function of k alone, $4\pi k^2 dk = d\mathbf{k}$ and this can be written

$$\alpha = - \int d\mathbf{k} k \partial_k C \gamma_k^{-1} \left[\frac{\gamma_k - \lambda k^2}{\gamma_k + \lambda k^2} f(\xi) + f(\eta) \right] . \quad (90)$$

To interpret the quantity $k \partial_k C$, we multiply the expression on the left hand side of (58), taken for $t_1 = t$, by $i k'_c$ to get

$$\begin{aligned} \left\langle \epsilon_{cde} \hat{z}_d^{+(0)}(\mathbf{k}) i k'_c z_e^{- (0)}(\mathbf{k}') \right\rangle &= - \left\langle \hat{z}_d^{+(0)}(\mathbf{k}) [i \mathbf{k}' \times \hat{\mathbf{z}}^{- (0)}(\mathbf{k}')]_d \right\rangle \\ &= - \left\langle \hat{\mathbf{z}}^{+(0)}(\mathbf{k}) \cdot \hat{\mathbf{w}}^{- (0)}(\mathbf{k}') \right\rangle , \end{aligned} \quad (91)$$

where

$$\mathbf{w}^{-(0)}(\mathbf{x}) \equiv \nabla \times \mathbf{z}^{-(0)}(\mathbf{x}) . \quad (92)$$

From Leslie (1973; eq. 2.16), (91) equals

$$- \hat{h}(\mathbf{k}) \delta(\mathbf{k} + \mathbf{k}') , \quad (93)$$

where $\hat{h}(\mathbf{k})$ is the transform of $h(\boldsymbol{\xi})$, the correlation function

$$h(\boldsymbol{\xi}) = \left\langle \mathbf{z}^{+(0)}(\mathbf{x}) \cdot \mathbf{w}^{-(0)}(\mathbf{x} + \boldsymbol{\xi}) \right\rangle . \quad (94)$$

From (61) with $t_1 = t$, we see that (91) is also equal to

$$\left\langle \epsilon_{cde} \hat{z}_d^{+(0)}(\mathbf{k}) i k'_c \hat{z}_e^{-(0)}(\mathbf{k}') \right\rangle = -i k_c (-2i k_c k^{-1} \partial_k C) \delta(\mathbf{k} + \mathbf{k}') = -2k \partial_k C \delta(\mathbf{k} + \mathbf{k}') . \quad (95)$$

Equating (93) with (95), we see that

$$k \partial_k C = \frac{1}{2} \hat{h}(\mathbf{k}) , \quad (96)$$

so that (90) becomes

$$\alpha = -\frac{1}{2} \int d\mathbf{k} \hat{h}(\mathbf{k}) \gamma_k^{-1} \left[\frac{\gamma_k - \lambda k^2}{\gamma_k + \lambda k^2} f(\xi) + f(\eta) \right] , \quad (97)$$

which is our principal result, where $\hat{h}(\mathbf{k})$, f , ξ , and η are defined by (94), (88), (82), and (84) respectively.

To compare (97) with the classical result (18), we take the limit $\overline{B} \rightarrow 0$, which according to (82) and (84) corresponds to $\xi \rightarrow 0$, $\eta \rightarrow 0$, in which case $f(\xi) = f(\eta) \rightarrow 1/3$, and the bracket in (97) becomes

$$\frac{1}{3} \left(\frac{\gamma_k - \lambda k^2}{\gamma_k + \lambda k^2} + 1 \right) = \frac{2}{3} \frac{\gamma_k}{\gamma_k + \lambda k^2} , \quad (98)$$

so

$$\alpha(\overline{B} \rightarrow 0) = -\frac{1}{3} \int d\mathbf{k} \hat{h}(\mathbf{k}) (\gamma_k + \lambda k^2)^{-1} . \quad (99)$$

As the integral of the power spectrum over $\mathbf{k}$ is the correlation at zero lag,

$$\begin{aligned}
\int d\mathbf{k} \hat{h}(\mathbf{k}) &= \langle \mathbf{z}^{+(0)}(\mathbf{x}, t) \cdot \nabla \times \mathbf{z}^{-(0)}(\mathbf{x}, t) \rangle \\
&= \langle (\mathbf{v}^{(0)} + \mathbf{b}^{(0)}) \cdot \nabla \times (\mathbf{v}^{(0)} - \mathbf{b}^{(0)}) \rangle \\
&= \langle \mathbf{v}^{(0)} \cdot \nabla \times \mathbf{v}^{(0)} \rangle - \langle \mathbf{b}^{(0)} \cdot \nabla \times \mathbf{b}^{(0)} \rangle + \langle \nabla \cdot (\mathbf{v}^{(0)} \times \mathbf{b}^{(0)}) \rangle \\
&= \langle \mathbf{v}^{(0)} \cdot \nabla \times \mathbf{v}^{(0)} \rangle - \langle \mathbf{b}^{(0)} \cdot \nabla \times \mathbf{b}^{(0)} \rangle
\end{aligned} \tag{100}$$

because $\langle \mathbf{v}^{(0)} \times \mathbf{b}^{(0)} \rangle = 0$ according to (2); here we have suppressed the arguments for clarity.

From (99) we conclude that if λ is small (but not too small — see below), and in the special case that $\gamma_k = \gamma_0$ is independent of k ,

$$\begin{aligned}
\alpha(\overline{B} \rightarrow 0) &= -\frac{1}{3} \int d\mathbf{k} \hat{h}(\mathbf{k}) \gamma_0^{-1} \\
&= -\frac{1}{3} \gamma_0^{-1} \left[\langle \mathbf{v}^{(0)}(\mathbf{x}, t) \cdot \nabla \times \mathbf{v}^{(0)}(\mathbf{x}, t) \rangle \right. \\
&\quad \left. - \langle \mathbf{b}^{(0)}(\mathbf{x}, t) \cdot \nabla \times \mathbf{b}^{(0)}(\mathbf{x}, t) \rangle \right] .
\end{aligned} \tag{101}$$

The first term agrees with classical result (18) if the damping constant γ_0 is identified with τ^{-1} . The second term has the sign and form expected from the work of Pouquet *et al.* (1976), as indicated in (19).

Our more general result for $\overline{B} \rightarrow 0$, (99), differs from the classical result in three ways:

- i) a term proportional to the current helicity is included in the form derived by Pouquet *et al.* (1976);
- ii) allowance is made for variation of the damping constant with k ;
- iii) and the effect of finite λ is included.

In the opposite limit $\overline{B} \rightarrow \infty$, $f(u) \rightarrow u^{-2}$, so the bracket in (97) becomes

$$\frac{\gamma_k - \lambda k^2}{\gamma_k + \lambda k^2} \frac{1}{u^2} + \frac{1}{v^2} = \frac{\gamma_k - \lambda k^2}{\gamma_k + \lambda k^2} \frac{(\gamma_k + \lambda k^2)^2}{(k\overline{B})^2} + \frac{(\lambda k^2)^2}{(k\overline{B})^2} = \left(\frac{\gamma_k}{k\overline{B}} \right)^2 , \tag{102}$$

independent of λ . Thus,

$$\alpha(\overline{B} \rightarrow \infty) = -\frac{1}{2\overline{B}^2} \int d\mathbf{k} \hat{h}(\mathbf{k}) k^{-2} \gamma_k. \quad (103)$$

An inverse dependence on $\overline{B}^2$ was cited by Krause and Rädler (1980), but with a different factor.

Finally, we consider the simplification introduced when the spectral density $C(k)$ is effectively concentrated at some wave number k_* , which if $k_* \sim 2k_0$ is a crude representation of Figure 1 of the numerical result of Pouquet *et al.* (1976), where E_k^V peaks at $k_0 \simeq 1$, E_k^M peaks at $k = 3$, and the energy beyond $k = 3$ is Alfvénic, so it does not contribute to α . Then from (82), (84), and (97),

$$\alpha = -\frac{1}{2} \int d\mathbf{k} \hat{h}(\mathbf{k}) \gamma_*^{-1} \left[\frac{\gamma_* - \lambda k_*^2}{\gamma_* + \lambda k_*^2} f(\xi_*) + f(\eta_*) \right], \quad (104)$$

where $\gamma_* = \gamma_k(k_*)$,

$$\xi_* = \frac{k_* \overline{B}}{\gamma_* + \lambda k_*^2}, \quad (105)$$

and

$$\eta_* = \frac{k_* \overline{B}}{\lambda k_*^2}. \quad (106)$$

In the light of (101), then,

$$\alpha = -\frac{1}{2} \gamma_*^{-1} \left[\langle \mathbf{v}^{(0)} \cdot \nabla \times \mathbf{v}^{(0)} \rangle - \langle \mathbf{b}^{(0)} \cdot \nabla \times \mathbf{b}^{(0)} \rangle \right] \left[\frac{\gamma_* - \lambda k_*^2}{\gamma_* + \lambda k_*^2} f(\xi_*) + f(\eta_*) \right], \quad (107)$$

so the dependence upon $\overline{B}$ is entirely in the multiplicative factor. If we define a magnetic Reynolds number by

$$R_M = \frac{\gamma_*}{\lambda k_*^2}, \quad (108)$$

and a nondimensional mean magnetic field by

$$\beta = \frac{\overline{B}}{(\gamma_*/k_*)}, \quad (109)$$

we can write the bracket in (107) in the form,

$$F(R_M, \beta) = \frac{R_M - 1}{R_M + 1} f\left(\beta \frac{R_M}{R_M + 1}\right) + f(\beta R_M). \quad (110)$$

As $\beta \rightarrow 0$, this approaches

$$F(R_M, \beta \rightarrow 0) = \frac{1}{3} \left(\frac{R_M - 1}{R_M + 1} + 1 \right) = \frac{2}{3} \frac{\gamma_*}{\gamma_* + \lambda k_*^2}, \quad (111)$$

which, when inserted into (107), yields agreement with (101), our previous result for $\overline{B} \rightarrow 0$.

However, one must be careful about this procedure if R_M is $\gg 1$, as is usually the case, for even if $\beta \ll 1$, β may be $\gg R_M^{-1}$, so that the second term on the right of (110) is effectively 0 rather than $1/3$. In this case

$$F(R_M \rightarrow \infty, \beta \rightarrow 0) = \frac{1}{3}, \quad (112)$$

and α is given by $1/2$ the value in (101). Evidently, the classical expression for α (as modified by Pouquet *et al.*) is off by a factor of 2 if $R_M \gg 1$. Figure 1 is a plot of $F(R_M, \beta)$, compared with $F^*(R_M, \beta) = (1 + R_M \beta^2)^{-1}$ from Cattaneo and Hughes (1996). As $F^*(R_M, \beta)$ was not proposed to be accurate for $R_M < 1$, the curves of F^* for $R_M < 1$ should be ignored. In qualitative agreement with $F(R_M, \beta)$, F^* decreases as R_M increases, but of critical importance, it decreases to 0, rather than remaining finite as $R_M \rightarrow \infty$, the case for the galactic dynamo.

4. Discussion

Our general result for α , (97), displays significant features of the α -effect due to driven helical turbulence. Our discussion of (101), above, makes clear that the values of $\mathbf{v}$ and $\mathbf{b}$ referred to in $\hat{h}(\mathbf{k})$ are $\mathbf{v}^{(0)}$ and $\mathbf{b}^{(0)}$, the values of the small-scale velocity and magnetic fields that apply if $\overline{\mathbf{B}} = 0$.

As indicated in the text, it is in principle possible to calculate $\mathbf{v}^{(0)}$ and $\mathbf{b}^{(0)}$, and hence $\hat{h}(\mathbf{k})$, from numerical simulations of driven helical turbulence with $\overline{\mathbf{B}} = 0$. Then evaluating α from (97) requires the value of γ_k , which can be calculated from (61) in terms of the statistics of the zero-order state. One should also calculate α directly from its definition using simulations with various values of $\overline{B}$, and then compare the results with (97).

Another feature of our work is that the contribution proportional to the current helicity, $\langle \mathbf{b}^{(0)} \cdot \nabla \times \mathbf{b}^{(0)} \rangle$, first described by Pouquet *et al.* (1976), emerges naturally from our work. To give it a physical interpretation, we refer to a derivation we carried out in terms of the variables $\mathbf{v}^{(0)}$ and $\mathbf{b}^{(0)}$ to check the derivation presented here in terms of $\mathbf{z}^{\pm(0)}$. Referring to (25), we find that at first order in $\overline{\mathbf{B}}$, the current helicity term is traceable to $\langle \mathbf{v}^{(1)} \times \mathbf{b}^{(0)} \rangle$, as opposed to $\langle \mathbf{v}^{(0)} \times \mathbf{b}^{(1)} \rangle$, which gives rise to the kinetic helicity term, where the velocity perturbation $\mathbf{v}^{(1)}$ is due to the first-order Lorentz force $(\nabla \times \mathbf{b}^{(0)}) \times \overline{\mathbf{B}}$ as explained by Pouquet *et al.* (1976; p. 332).

Current helicity is discussed in the Soviet and Russian literature. Vainshtein (1972) and Vainshtein and Zeldovich (1972) called attention to effects of current helicity (which they referred to as “magnetic gyrotropy”). They argued on physical grounds that as $\overline{B}$ grows, it induces current helicity, which, because its sign is opposite to that of $\langle \mathbf{v} \cdot \nabla \times \mathbf{v} \rangle$, reduces α , causing dynamo activity to cease at some finite value of $\overline{B}$. As we have explained, Pouquet *et al.* (1976) showed that current helicity is important at all values of $\overline{B}$, and on the basis of their spectral equations, found that the effect described by Vainshtein (1972) would be smaller and of the opposite sign than he claimed. Vainshtein and Kichaninov (1983) accept the interpretation of Pouquet *et al.* (1976). They also derive a differential equation for the small-scale current helicity from the conservation of magnetic helicity as did Kleeorin and Ruzmaikin *et al.* (1982); see Kleeorin *et al.* (1995). We discuss these matters further in a forthcoming paper (Blackman and Field 1998).

The present work relies upon $\tau \ll t_{ed}$ to evaluate the perturbations of $\mathbf{v}$ and $\mathbf{b}$ due to $\overline{\mathbf{B}}$ (although not in evaluating $\mathbf{v}^{(0)}$ and $\mathbf{b}^{(0)}$). This is a reasonable approximation because the dominant contribution to α comes from wavenumbers $< 3k_0$, which are relatively unaffected by magnetic back reaction, and therefore should follow Pope’s (1994) finding that in hydrodynamic turbulence, $\tau < t_{ed}$. Unfortunately, the inequality is not as strong as we desire.

5. Conclusions

We have found an analytic formula for the dynamo coefficient α based upon a perturbation expansion in $\overline{B}$, the magnitude of the large-scale magnetic field, which is responsible for perturbing isotropic helical turbulence in such a way as to produce a turbulent emf along $\overline{\mathbf{B}}$. Our formula gives α in terms of $\overline{B}$ and the magnetic diffusivity λ , together with the spectra of kinetic and current helicities, and a damping coefficient. Both of the latter are calculable from a simulation of incompressible isotropic helical MHD turbulence. For small values of $\overline{B}$, our results agree with the classical results as modified by Pouquet *et al.* (1976). For large $\overline{B}$, we find that two terms contribute to α . The first term is independent of the magnetic diffusivity λ , and is similar to the expression proposed by Kraichnan (1979). The second term, which depends on λ in a way reminiscent of the expressions suggested by Cattaneo and his collaborators, vanishes in the limit $\lambda \rightarrow 0$, as predicted by Cattaneo, *et al.*, but contrary to Cattaneo *et al.*, the first term remains finite, and so α is reduced only by a factor of 2 for large values of $\overline{B}$.

We are grateful to Amitava Bhattacharjee, B. D. G. Chandran, and Eugene Parker for their insightful comments, and to the Aspen Center for Physics, where part of the work was done. We are also grateful to Mr. Dimitri Kalaitzidakis for his hospitality and allowing us

to use his facilities while at a meeting on Crete. This work was supported in part by NASA Grant NAGW-931.

A. Energy Cascade

Start with the Navier-Stokes equation, which describes incompressible hydrodynamic (HD) turbulence:

$$\partial_t \mathbf{v} = -\mathbf{v} \cdot \nabla \mathbf{v} - \nabla p + \nu \nabla^2 \mathbf{v} + \mathbf{f} , \quad (\text{A1})$$

where $p = \text{pressure} \div \rho$. Define the kinetic energy E^V by

$$E^V = \left\langle \frac{1}{2} v^2 \right\rangle . \quad (\text{A2})$$

Then the scalar product of (A1) with $\mathbf{v}$ shows that

$$\partial_t E^V = -\nu \left\langle \omega^2 \right\rangle + \langle \mathbf{f} \cdot \mathbf{v} \rangle , \quad (\text{A3})$$

where

$$\boldsymbol{\omega} = \nabla \times \mathbf{v} \quad (\text{A4})$$

is the vorticity. Hence in unforced ideal HD, E^V is conserved. The only effect of the nonlinear term $-\mathbf{v} \cdot \nabla \mathbf{v}$ in (A.1) is to redistribute E_k^V , the kinetic energy spectrum, over k , keeping $E^V = \int E_k^V dk$ constant. Numerical simulations show that this redistribution takes the form of a direct cascade, that is, from lower k to higher k .

Now add a magnetic field $\mathbf{B}$, so the momentum equation becomes (25):

$$\begin{aligned} \partial_t \mathbf{v} &= -\mathbf{v} \cdot \nabla \mathbf{v} - \nabla p + \mathbf{B} \cdot \nabla \mathbf{B} - \nabla \frac{1}{2} B^2 + \nu \nabla^2 \mathbf{v} + \mathbf{f} \\ &= -\mathbf{v} \cdot \nabla \mathbf{v} + \mathbf{B} \cdot \nabla \mathbf{B} - \nabla P + \nu \nabla^2 \mathbf{v} + \mathbf{f} , \end{aligned} \quad (\text{A5})$$

where

$$P = p + \frac{1}{2} B^2 . \quad (\text{A6})$$

The equation for E^V becomes

$$\partial_t E^V = \langle \mathbf{v} \cdot (\mathbf{B} \cdot \nabla) \mathbf{B} \rangle - \nu \langle \omega^2 \rangle + \langle \mathbf{f} \cdot \mathbf{v} \rangle , \quad (\text{A7})$$

so E^V is not conserved in unforced ideal MHD because the first term allows magnetic energy to exchange with kinetic energy. To account for this, we write the induction equation governing $\mathbf{B}$:

$$\partial_t \mathbf{B} = -\mathbf{v} \cdot \nabla \mathbf{B} + \mathbf{B} \cdot \nabla \mathbf{v} + \nu \nabla^2 \mathbf{B} , \quad (\text{A8})$$

where we have specialized to the case of unit magnetic Prandtl number, λ/ν . From the scalar product of (A.8) with $\mathbf{B}$ we obtain an equation for the magnetic energy $E^M = \langle \frac{1}{2} B^2 \rangle$,

$$\partial_t E^M = \langle \mathbf{B} \cdot (\mathbf{B} \cdot \nabla) \mathbf{v} \rangle - \nu \langle J^2 \rangle , \quad (\text{A9})$$

where

$$\mathbf{J} = \nabla \times \mathbf{B} . \quad (\text{A10})$$

E^M is not conserved in unforced ideal MHD because of exchange with E^V . However, we note that in (A.10)

$$\begin{aligned} \langle \mathbf{B} \cdot (\mathbf{B} \cdot \nabla) \mathbf{v} \rangle &= \langle \mathbf{B} \cdot \nabla (\mathbf{v} \cdot \mathbf{B}) \rangle - \langle \mathbf{v} \cdot (\mathbf{B} \cdot \nabla) \mathbf{B} \rangle \\ &= \langle \nabla \cdot [\mathbf{B} (\mathbf{v} \cdot \mathbf{B})] \rangle - \langle \mathbf{v} \cdot (\mathbf{B} \cdot \nabla) \mathbf{B} \rangle \\ &= - \langle \mathbf{v} \cdot (\mathbf{B} \cdot \nabla) \mathbf{B} \rangle \end{aligned} \quad (\text{A11})$$

since spatial averaging converts the divergence to a vanishing surface integral. Hence

$$\partial_t E^M = - \langle \mathbf{v} \cdot (\mathbf{B} \cdot \nabla) \mathbf{B} \rangle - \nu \langle J^2 \rangle . \quad (\text{A12})$$

The energy-exchange term in (A12) is the negative of that in (A7), indicating that energy gained by E^V is lost by E^M and *vice-versa*. Hence if $\nu = \mathbf{f} = 0$, the sum of (A.7) and (A.13) yields

$$\partial_t E \equiv \partial_t (E^V + E^M) = 0 \quad (\text{A13})$$

for the conservation of the total energy E in unforced ideal MHD. By analogy with E in HD, the second nonlinear terms in (A.5) and (A.8) for MHD can only redistribute E_k over k . Numerical studies like those of Pouquet *et al.* (1976) show that in the non helical case, the effect of the nonlinear terms in MHD is a direct cascade of total energy. However, because E^V and E^M are not separately conserved, it is not obvious that both E^V and E^M individually cascade directly in general.

To answer this question, we turn to the Elssässer variables for the small-scale field $\mathbf{z}^\pm$, which in the case $\lambda = \nu$ satisfy (35):

$$\partial_t \mathbf{z}^\pm = -\mathbf{z}^\mp \cdot \nabla \mathbf{z}^\pm - \nabla P + \nu \nabla^2 \mathbf{z}^\pm + \mathbf{f} \pm \bar{\mathbf{B}} \cdot \nabla \mathbf{z}^\pm . \quad (\text{A14})$$

Define

$$E^\pm = \left\langle \frac{1}{2} (z^\pm)^2 \right\rangle . \quad (\text{A15})$$

Then the scalar product of (A.15) with $\mathbf{z}^\pm$ yields

$$\begin{aligned} \partial_t E^\pm &= -\left\langle \mathbf{z}^\pm \cdot (\mathbf{z}^\mp \cdot \nabla) \mathbf{z}^\pm \right\rangle - \left\langle \mathbf{z}^\pm \cdot \nabla P \right\rangle \\ &\quad + \nu \left\langle \mathbf{z}^\pm \cdot \nabla^2 \mathbf{z}^\pm \right\rangle + \left\langle \mathbf{f} \cdot \mathbf{z}^\pm \right\rangle \pm \left\langle \mathbf{z}^\pm \cdot (\bar{\mathbf{B}} \cdot \nabla) \mathbf{z}^\pm \right\rangle \\ &= -\left\langle (\mathbf{z}^\mp \cdot \nabla) \frac{1}{2} (z^\pm)^2 \right\rangle - \left\langle \nabla \cdot (\mathbf{z}^\pm P) \right\rangle \\ &\quad + \nu \left\langle \mathbf{z}^\pm \cdot [\nabla(\nabla \cdot \mathbf{z}^\pm) - \nabla \times \nabla \times \mathbf{z}^\pm] \right\rangle \\ &\quad + \left\langle \mathbf{f} \cdot \mathbf{z}^\pm \right\rangle \pm \left\langle \bar{\mathbf{B}} \cdot \nabla \frac{1}{2} (z^\pm)^2 \right\rangle . \end{aligned} \quad (\text{A16})$$

Since $\nabla \cdot \mathbf{z}^\pm = 0$, this gives

$$\begin{aligned} \partial_t E^\pm &= -\left\langle \nabla \cdot \left[\frac{1}{2} \mathbf{z}^\mp (z^\pm)^2 \right] \right\rangle - \left\langle \nabla \cdot (\mathbf{z}^\pm P) \right\rangle \\ &\quad - \nu \left\langle \mathbf{z}^\pm \cdot \nabla \times \boldsymbol{\omega}^\pm \right\rangle + \left\langle \mathbf{f} \cdot \mathbf{z}^\pm \right\rangle \pm \left\langle \nabla \cdot [\bar{\mathbf{B}} \frac{1}{2} (z^\pm)^2] \right\rangle , \end{aligned} \quad (\text{A17})$$

where

$$\boldsymbol{\omega}^\pm = \nabla \times \mathbf{z}^\pm = \boldsymbol{\omega} \pm \mathbf{j} . \quad (\text{A18})$$

Since

$$\begin{aligned} \mathbf{z}^\pm \cdot \nabla \times \boldsymbol{\omega}^\pm &= \nabla \cdot (\boldsymbol{\omega}^\pm \times \mathbf{z}^\pm) + \boldsymbol{\omega}^\pm \cdot \nabla \times \mathbf{z}^\pm \\ &= \nabla \cdot (\boldsymbol{\omega}^\pm \times \mathbf{z}^\pm) + (\boldsymbol{\omega}^\pm)^2 \end{aligned} \quad (\text{A19})$$

and averages of divergences vanish,

$$\partial_t E^\pm = -\nu \langle (\boldsymbol{\omega}^\pm)^2 \rangle + \langle \mathbf{f} \cdot \mathbf{z}^\pm \rangle, \quad (\text{A20})$$

an equation identical in form to (A.3). We conclude that if $\lambda = \nu$ both E^+ and E^- are individually conserved in unforced ideal MHD, and that the effect of nonlinear interactions is to redistribute the two energy spectra E_k^+ and E_k^- individually. Note that the presence or absence of a mean magnetic field $\overline{\mathbf{B}}$ does not affect this conclusion.

That leaves the question as to whether the redistribution is a direct cascade. Since

$$\begin{aligned} E^\pm &= \left\langle \frac{1}{2} (z^\pm)^2 \right\rangle = \left\langle \frac{1}{2} (\mathbf{v} \pm \mathbf{B})^2 \right\rangle \\ &= \left\langle \frac{1}{2} v^2 \right\rangle + \left\langle \frac{1}{2} B^2 \right\rangle \pm \langle \mathbf{v} \cdot \mathbf{B} \rangle \\ &= E^V + E^M \pm 2K = E \pm 2K, \end{aligned} \quad (\text{A21})$$

where

$$K = \frac{1}{2} \langle \mathbf{v} \cdot \mathbf{B} \rangle \quad (\text{A22})$$

is the cross helicity (Biskamp 1993, p. 179), it follows from the conservation of E^+ , E^- , and E that K is also conserved in unforced ideal MHD if $\lambda = \nu$. As pointed out by Pouquet *et al.* (1976), if the initial state is assumed to be statistically invariant under $\mathbf{b} \rightarrow -\mathbf{b}$, it vanishes initially and thus remains equal to zero, and from (A21) we have

$$E^+ = E^- = E. \quad (\text{A23})$$

As E cascades directly, it follows that E^+ and E^- cascade directly if $\lambda = \nu$.

B. Correlations

Here we show that (58) is a reasonable representation. Let

$$\mathbf{V} = \left\langle \hat{\mathbf{z}}^{+(0)}(\mathbf{k}, t) \times \mathbf{z}^{-(0)}(\mathbf{k}', t_1) \right\rangle , \quad (\text{B1})$$

so that the average in (57) is V_c . From (50)

$$\begin{aligned} \mathbf{V} &= (2\pi)^{-6} \int \int d\mathbf{x} d\mathbf{x}' e^{-i(\mathbf{k} \cdot \mathbf{x} + \mathbf{k}' \cdot \mathbf{x}')} \left\langle \mathbf{z}^{+(0)}(\mathbf{x}, 0) \times \mathbf{z}^{-(0)}(\mathbf{x}', \tau) \right\rangle \\ &= (2\pi)^{-6} \int \int d\mathbf{x} d\mathbf{r} e^{-i(\mathbf{k} + \mathbf{k}') \cdot \mathbf{x} - i\mathbf{k}' \cdot \mathbf{r}} \left\langle \mathbf{z}^{+(0)}(0, 0) \times \mathbf{z}^{-(0)}(\mathbf{r}, \tau) \right\rangle , \end{aligned} \quad (\text{B2})$$

where we have put $t_1 = t + \tau$ and $\mathbf{x}' = \mathbf{x} + \mathbf{r}$ and have used stationarity and homogeneity.

Carrying out the integration over $\mathbf{x}$ gives

$$\mathbf{V} = (2\pi)^{-3} \delta(\mathbf{k} + \mathbf{k}') \hat{\mathbf{I}}(\mathbf{k}', \tau) , \quad (\text{B3})$$

where

$$\hat{\mathbf{I}}(\mathbf{k}', \tau) = \int d\mathbf{r} e^{-i\mathbf{k}' \cdot \mathbf{r}} \left\langle \mathbf{z}^{+(0)}(0, 0) \times \mathbf{z}^{-(0)}(\mathbf{r}, \tau) \right\rangle . \quad (\text{B4})$$

If we write

$$\mathbf{Q}(\mathbf{r}, \tau) = \left\langle \mathbf{v}(0, 0) \times \mathbf{v}(\mathbf{r}, \tau) \right\rangle , \quad (\text{B5})$$

$$\mathbf{S}(\mathbf{r}, \tau) = \left\langle \mathbf{b}(0, 0) \times \mathbf{b}(\mathbf{r}, \tau) \right\rangle , \quad (\text{B6})$$

$$\mathbf{P}_v(\mathbf{r}, \tau) = \left\langle \mathbf{v}(0, 0) \times \mathbf{b}(\mathbf{r}, \tau) \right\rangle , \quad (\text{B7})$$

and

$$\mathbf{P}_b(\mathbf{v}, \tau) = \left\langle \mathbf{b}(0, 0) \times \mathbf{v}(\mathbf{r}, \tau) \right\rangle , \quad (\text{B8})$$

where we have omitted superscripts for clarity, then the average in (B4) is, from (20) and (21),

$$\begin{aligned} \left\langle \mathbf{z}^{+(0)}(0, 0) \times \mathbf{z}^{-(0)}(\mathbf{r}, \tau) \right\rangle &= \mathbf{Q}(\mathbf{r}, \tau) - \mathbf{S}(\mathbf{r}, \tau) \\ &\quad - \mathbf{P}_v(\mathbf{r}, \tau) + \mathbf{P}_b(\mathbf{r}, \tau) , \end{aligned} \quad (\text{B9})$$

so

$$\hat{\mathbf{I}} = \hat{\mathbf{Q}}(\mathbf{k}', \tau) - \hat{\mathbf{S}}(\mathbf{k}', \tau) - \hat{\mathbf{P}}_v(\mathbf{k}', \tau) + \hat{\mathbf{P}}_b(\mathbf{k}', \tau) , \quad (\text{B10})$$

where

$$\hat{\mathbf{Q}}(\mathbf{k}', \tau) = \int d\mathbf{r} e^{-i\mathbf{k}' \cdot \mathbf{r}} \mathbf{Q}(\mathbf{r}, \tau) , \quad (\text{B11})$$

etc. One of us has shown (Chou & Fish 1998) that the terms $\mathbf{P}_v$ and $\mathbf{P}_b$ do not contribute to the final result, so we ignore them in what follows.

Consider

$$\begin{aligned} \mathbf{Q}(\mathbf{r}, \tau) &= \langle \mathbf{v}(0, 0) \times \mathbf{v}(\mathbf{r}, \tau) \rangle \\ &= \langle \mathbf{v}(-\mathbf{r}, -\tau) \times \mathbf{v}(0, 0) \rangle \\ &= - \langle \mathbf{v}(0, 0) \times \mathbf{v}(-\mathbf{r}, -\tau) \rangle , \end{aligned} \quad (\text{B12})$$

where we have used homogeneity and stationarity and flipped the cross product. Thus

$$\begin{aligned} \hat{\mathbf{Q}}(\mathbf{k}', \tau) &= - \int d\mathbf{r} e^{-i\mathbf{k}' \cdot \mathbf{r}} \langle \mathbf{v}(0, 0) \times \mathbf{v}(-\mathbf{r}, -\tau) \rangle \\ &= - \int d\mathbf{r} e^{i\mathbf{k}' \cdot \mathbf{p}} \langle \mathbf{v}(0, 0) \times \mathbf{v}(\mathbf{p}, -\tau) \rangle \\ &= -\hat{\mathbf{Q}}(-\mathbf{k}', -\tau) . \end{aligned} \quad (\text{B13})$$

Krause and Rädler (1980, p. 75) show that quantities like $\mathbf{Q}$ are odd functions of $\mathbf{k}'$, so (B13) implies that

$$\hat{\mathbf{Q}}(\mathbf{k}', \tau) = \hat{\mathbf{Q}}(\mathbf{k}', -\tau) . \quad (\text{B14})$$

Since $\mathbf{S}(\mathbf{r}, \tau)$ has a similar structure,

$$\hat{\mathbf{S}}(\mathbf{k}', \tau) = \hat{\mathbf{S}}(\mathbf{k}', -\tau) , \quad (\text{B15})$$

so that $\hat{\mathbf{I}}(\mathbf{k}', \tau)$ is an even function of τ , or, equivalently, a function of $|\tau|$, which we shall denote $(2\pi)^3 \mathbf{R}(\mathbf{k}', |\tau|)$. Hence from (B3)

$$V_c = \delta(\mathbf{k} + \mathbf{k}') R_c(\mathbf{k}', |\tau|) . \quad (\text{B16})$$

Because R is composed of cross products, we introduce a tensor $R_{de}(\mathbf{k}', |\tau|)$ such that

$$R_c(\mathbf{k}', |\tau|) = \epsilon_{cde} R_{de}(\mathbf{k}', |\tau|) , \quad (\text{B17})$$

so that

$$V_c = \epsilon_{cde} \delta(\mathbf{k} + \mathbf{k}') R_{de}(\mathbf{k}', |\tau|) . \quad (\text{B18})$$

The last step is to assume that $R_{de}(\mathbf{k}', |\tau|)$ is a function of $\mathbf{k}'$ times a function of $|\tau|$ (which may depend upon $k = |\mathbf{k}| = |\mathbf{k}'|$):

$$R_{de}(\mathbf{k}', |\tau|) = R_{de}(\mathbf{k}') f_k(|\tau|) , \quad (\text{B19})$$

where by taking $f_k(0) = 1$, we assure that $R_{de}(\mathbf{k}')$ represents the maximum value of the correlation. As a natural choice for $f_k(|\tau|)$ we take

$$f_k(|\tau|) = e^{-\gamma_k |\tau|} , \quad (\text{B20})$$

where γ_k is a k -dependent inverse correlation time, in the spirit of the eddy-damping approximation of Pouquet et al. (1976) and Kraichnan (1979). Then

$$V_c = \epsilon_{cde} \delta(\mathbf{k} + \mathbf{k}') R_{de}(\mathbf{k}') e^{-\gamma_k |\tau|} , \quad (\text{B21})$$

which is (58).

REFERENCES

- Biskamp, D. 1993, *Nonlinear Magnetohydrodynamics*, (Cambridge: Cambridge Univ. Press)
- Blackman, E. G. & Field, G. B. 1998, in preparation
- Cattaneo, F. 1994, *ApJ*, 434, 200
- Cattaneo, F. & Hughes, D. W. 1996, *Phys. Rev. E.*, 54, 4532
- Cattaneo, F. & Vainshtein, S.I. 1991, *ApJ*, 376, L21
- Chandran, B. D. G. 1997, *ApJ*, 485, 148
- Chou, H. & Fish, V. 1998, submitted to *ApJ*
- Frisch, U., Pouquet, A., L  orat, J. & Mazure, A. 1975, *J. Fluid Mech.*, 68, pt. 4, 769
- Keinigs, R. K. 1983, *Phys. Fluids*, 26, 2558
- Kichanitov, L. L. 1985, *Magn. Hidro.* 2, 3
- Kleeorin, N. & Ruzmaikin, A., 1982, *Mag. Hidro.* N2, 17
- Kleeorin, N., Rogachevskii, I., & Ruzmaikin, A. 1995, *A&A*, 297, 159
- Kraichnan, R. H. 1979, *Phys. Rev. Lett.*, 42, 1677
- Krause, F. & R  dler, K.H. 1980, *Mean-Field Magnetohydrodynamics and Dynamo Theory*,
(Oxford: Pergamon)
- Kulsrud, R.M. & Anderson, S.W. 1992, *ApJ*, 396, 606
- Leslie, D. C. 1973, *Developments in the Theory of Turbulence*, (Oxford, Clarendon Press)
- Meneguzzi, M., Frisch, U., & Pouquet, A. 1981, *Phys. Rev. Lett.*, 47, 1060

- Moffatt, H.K. 1978, *Magnetic Field Generation in Electrically Conducting Fluids*,
(Cambridge: Cambridge University Press)
- Montgomery, D. & Chen, H. 1984, *Plasma Phys. & Controlled Fusion*, 26, 1189
- Parker, E. N. 1955, *ApJ*, 122, 293
- Parker, E. N. 1971, *ApJ*, 163, 255
- Parker, E. N. 1979, *Cosmical Magnetic Fields*, (Oxford: Clarendon Press)
- Piddington, J. H. 1970, *Aust. J. Phys.*, 23, 731
- Piddington, J. H. 1972a, *Sol. Phys.*, 22, 3
- Piddington, J. H. 1972b, *Cos. Elec.*, 3, 50
- Piddington, J. H. 1972c, *ibid.*, 5, 129
- Piddington, J. H. 1975a, *Ap. & Sp. Sci.*, 35, 269
- Piddington, J. H. 1975b, *ibid.*, 39, 157
- Pope, S. B. 1994, *Ann. Rev. Fl. Mech.*, 26, 23
- Pouquet, A., Frisch, U., & Léorat, J. 1976, *J. Fl. Mech.*, 77, pt. 2, 321
- Pouquet, A. & Patterson, G. S. 1978, *J. Fl. Mech.*, 85, pt. 2, 305
- Ruzmaikin, A.A., Shukurov, A.M. & Sokoloff, D.D. 1988, *Magnetic Fields of Galaxies*,
(Dordrecht: Kluwer)
- Seehafer, N. 1994, *Europhys. Lett.*, 27, 353
- Seehafer, N. 1995, *Astron. & Astrophys.*, 301, 290

- Tao, L., Cattaneo, F., & Vainshtein, S.I. 1993, in Theory of Solar and Planetary Dynamos, ed. M.R.E. Proctor, P.C. Matthews & A.M. Rucklidge, p.303, (Cambridge: Cambridge University Press)
- Vainshtein, S. I. 1972, Sov. Phys. JETP, 34, 327
- Vainshtein, S. I. & Cattaneo, F. 1992, ApJ, 393, 165
- Vainshtein, S. I. & Kichanitov, L. L. 1983, Geophys. Astrophys. Fluid Dynamics, 24, 273
- Vainshtein, S. I. & Rosner, R. 1991, ApJ, 376, 199
- Vainshtein, S. I. & Zeldovich, Y. B. 1972, Sov. Phys. Uspekhi, 15, 159

Fig. 1.— Three functions of the magnetic Reynolds number R_M and the normalized magnetic field $\beta = \overline{B}/v_0$ are plotted and compared. $F(R_M, \beta)$, defined in (110) and represented by solid lines, is our result for the dependence of α on R_M for $\beta = 10^{-5}$, 3×10^{-3} , and 1. $F(R_M, \beta \rightarrow 0)$, defined in (109) and represented by the circles, is our result for $\beta \rightarrow 0$. $F^*(R_M, \beta) = (1 + R_M \beta^2)^{-1}$ is the dependence of α suggested by Cattaneo & Hughes (1996); as it was not intended to be accurate for $R_M < 1$, points for $R_M < 1$ should be ignored.

